

Finite groups determined by an inequality of the orders of their subgroups II

Marius Tărnăuceanu

October 25, 2016

Abstract

In this note we study a class of finite groups for which the orders of subgroups satisfy a certain inequality. In particular, characterizations of the well-known groups $\mathbb{Z}_2 \times \mathbb{Z}_2$ and S_3 are obtained.

MSC2000: Primary 20D60, 20D30; Secondary 20D15, 11A25.

Key words: finite groups, subgroup lattices, number of subgroups, arithmetic functions.

1 Introduction

Let G be a finite group, $L(G)$ be the subgroup lattice of G and

$$\sigma_1(G) = \sum_{H \in L(G)} \frac{|H|}{|G|} = \sum_{H \in L(G)} \frac{1}{|G:H|}.$$

The starting point for our discussion is given by the paper [1], where the groups G with $\sigma_1(G) \leq 2$ have been determined. Recall also several basic properties of the function σ_1 :

- if G is cyclic of order n and $\sigma(n)$ denotes the sum of all divisors of n , then $\sigma_1(G) = \frac{\sigma(n)}{n}$;
- σ_1 is multiplicative, i.e. if G_i , $i = 1, 2, \dots, m$, are finite groups of coprime orders, then $\sigma_1(\prod_{i=1}^m G_i) = \prod_{i=1}^m \sigma_1(G_i)$;

$$- \sigma_1(G) \geq \sigma_1(G/H), \forall H \trianglelefteq G.$$

By refining the proof of Theorem 1 of [1], in the current paper we will determine the finite groups G satisfying $\sigma_1(G) \leq 2 + \frac{4}{|G|}$. Our main result is the following.

Theorem 1. *Let G be a finite group of order n . Then:*

- a) $\sigma_1(G) < 2 + \frac{4}{n}$ if and only if G is cyclic and $\sigma(n) < 2n + 4$ or $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2$;
- b) $\sigma_1(G) = 2 + \frac{4}{n}$ if and only if G is cyclic and $\sigma(n) = 2n + 4$ or $G \cong \mathbb{Z}_3 \times \mathbb{Z}_3$ or $G \cong S_3$.

Two nice characterizations of $\mathbb{Z}_2 \times \mathbb{Z}_2$ and S_3 can be inferred from the above theorem.

Corollary 2. $\mathbb{Z}_2 \times \mathbb{Z}_2$ is the unique non-cyclic group G satisfying $\sigma_1(G) < 2 + \frac{4}{|G|}$, while S_3 is the unique non-abelian group G satisfying $\sigma_1(G) = 2 + \frac{4}{|G|}$.

By Theorem 1 of [1], a finite group G with $\sigma_1(G) \leq 2$ is cyclic of deficient or perfect order. Also, in Lemma 4 below we will show that a finite group G with $\sigma_1(G) < 2 + \frac{4}{|G|}$ is always nilpotent. Inspired by these results, we came up with the following natural problem: is there a constant $c \in (2, \infty)$ such that if $\sigma_1(G) < c$ then G is nilpotent? The answer to this problem is negative, as shows our next theorem.

Theorem 3. *There are sequences of finite non-nilpotent groups $(G_n)_{n \in \mathbb{N}}$ such that $\sigma_1(G_n) \searrow 2$ for $n \rightarrow \infty$.*

Finally, we note that an interesting open problem is whether there is a constant $c \in (2, \infty)$ such that if $\sigma_1(G) < c$ then G is solvable.

Most of our notation is standard and will not be repeated here. Basic definitions and results on groups can be found in [3]. For subgroup lattice concepts we refer the reader to [2] and [4].

2 Proof of the main results

We start by proving two auxiliary results.

Lemma 4. *Let G be a finite group. If $\sigma_1(G) < 2 + \frac{4}{|G|}$ then G is nilpotent, while if $\sigma_1(G) = 2 + \frac{4}{|G|}$ then G is nilpotent or $G \cong S_3$.*

Proof. Assume that $\sigma_1(G) < 2 + \frac{4}{|G|}$ and G is not nilpotent, that is it contains a non-normal maximal subgroup M . Then M coincides with its normalizer in G and so it has exactly $r = |G : M|$ conjugates, say $M_1, M_2, \dots, M_r$. On the other hand, $L(G)$ cannot consist only of $1, G$ and $M_1, M_2, \dots, M_r$. Therefore there is $N \leq G$ such that $N \neq 1, G, M_1, M_2, \dots, M_r$. Since

$$2 + \frac{4}{|G|} > \sigma_1(G) \geq \frac{1 + r|M| + |N| + |G|}{|G|} = 2 + \frac{|N| + 1}{|G|},$$

it follows that N is a normal subgroup of order 2 and $L(G) = \{1, G, M_1, M_2, \dots, M_r, N\}$. Then either $N \subset M_i, \forall i = 1, 2, \dots, r$, or $N \cap M_i = 1, \forall i = 1, 2, \dots, r$. We infer that G is either a 2-group or a cyclic group of order $2p$ for some odd prime p , a contradiction.

Assume now that $\sigma_1(G) = 2 + \frac{4}{|G|}$ and G is not nilpotent. Then, under the above notation, we must have $|N| = 3$ and thus G is a group of order $3q$ for some prime q . Clearly, the conditions $r \mid 3$ and $r \equiv 1 \pmod{q}$ imply $r = 3$ and $q = 2$. Hence $G \cong S_3$, completing the proof. $\square$

Lemma 5. *Let G be a non-cyclic p -group. If $\sigma_1(G) < 2 + \frac{4}{|G|}$ then $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2$, while if $\sigma_1(G) = 2 + \frac{4}{|G|}$ then $G \cong \mathbb{Z}_3 \times \mathbb{Z}_3$.*

Proof. Let $|G| = p^n$ and $G/\Phi(G) \cong \mathbb{Z}_p^k$. Then $2 \leq k \leq n$. Assume that $\sigma_1(G) \leq 2 + \frac{4}{p^n}$ and denote by $a_{k,p}(i)$ the number of subgroups of order p^i in $\mathbb{Z}_p^k$, $i = 0, 1, \dots, k$. If $k \geq 3$ then

$$\begin{aligned} \sigma_1(G) &\geq \frac{1}{p^n} (a_{k,p}(0)p^{n-k} + a_{k,p}(1)p^{n-k+1} + a_{k,p}(k-1)p^{n-1} + a_{k,p}(k)p^n) \\ &= \frac{1}{p^k} (a_{k,p}(0) + a_{k,p}(1)p + a_{k,p}(k-1)p^{k-1} + a_{k,p}(k)p^k) \\ &= \frac{1}{p^k} \left(1 + \frac{p^k - 1}{p - 1}p + \frac{p^k - 1}{p - 1}p^{k-1} + p^k \right) \\ &> \frac{1}{p^k} (1 + 4p^k) > 4 > 2 + \frac{4}{p^n}, \end{aligned}$$

contradicting our hypothesis. Consequently, $k = 2$. For $n > 2$ we infer that

$$2 + \frac{4}{p^n} \geq \sigma_1(G) \geq \frac{1 + (p+1)p^{n-1} + p^{n-2} + p^n}{p^n},$$

which leads to

$$4 \geq p^{n-1} + p^{n-2} + 1,$$

a contradiction. Therefore $n = 2$, i.e. $G \cong \mathbb{Z}_p \times \mathbb{Z}_p$. Moreover, we have

$$2 + \frac{4}{p^2} \geq \sigma_1(G) = \frac{1 + (p+1)p + p^2}{p^2} \iff 4 \geq p+1 \iff p \in \{2, 3\}.$$

Thus $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ for $\sigma_1(G) < 2 + \frac{4}{p^2}$ and $G \cong \mathbb{Z}_3 \times \mathbb{Z}_3$ for $\sigma_1(G) = 2 + \frac{4}{p^2}$, as desired. $\square$

We are now able to prove our main results.

Proof of Theorem 1. a) Assume that $\sigma_1(G) < 2 + \frac{4}{n}$. Then G is nilpotent by Lemma 4, which implies that it can be written as a direct product of its Sylow p_i -subgroups:

$$G \cong \prod_{i=1}^m G_i. \quad (*)$$

For every $i = 1, 2, \dots, m$, we have

$$\sigma_1(G_i) \leq \sigma_1(G) < 2 + \frac{4}{n} \leq 2 + \frac{4}{|G_i|}.$$

By Lemma 5 it follows that either G_i is cyclic or $G_i \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. We infer that either G is cyclic or $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_{n'}$ for some odd positive integer n' . Suppose that G is not cyclic. If $n' > 1$ we obtain

$$2 + \frac{1}{n'} = 2 + \frac{4}{n} > \sigma_1(G) = \sigma_1(\mathbb{Z}_2 \times \mathbb{Z}_2) \sigma_1(\mathbb{Z}_{n'}) = \frac{11\sigma(n')}{4n'},$$

or equivalently

$$11\sigma(n') < 8n' + 4,$$

a contradiction. Thus $n' = 1$ and $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.

b) Assume that $\sigma_1(G) = 2 + \frac{4}{n}$ and $G \not\cong S_3$. Then G is nilpotent by Lemma 4 and so it has a direct decomposition of type (*). For $m \geq 2$ we have

$$\sigma_1(G_i) \leq \sigma_1(G) = 2 + \frac{4}{n} < 2 + \frac{4}{|G_i|}, \quad i = 1, 2, \dots, m,$$

and again every G_i is cyclic or isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$ by Lemma 5. Therefore either G is cyclic or $G \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_{n'}$ for some odd positive integer n' . In the second case the condition $\sigma_1(G) = 2 + \frac{4}{n}$ leads to

$$11\sigma(n') = 8n' + 4,$$

a contradiction. For $m = 1$ it follows that G is a p -group, and consequently it is cyclic or isomorphic to $\mathbb{Z}_3 \times \mathbb{Z}_3$ by Lemma 5. This completes the proof. $\square$

Proof of Theorem 3. Let $(p_n)_{n \in \mathbb{N}}$ be the sequence of prime numbers. By Dirichlet's theorem we infer that for every $n \in \mathbb{N}$ there is a prime q_n such that $p_n \mid q_n - 1$. Let G_n be the non-nilpotent group of order $p_n q_n$. This contains one subgroup of order 1, q_n subgroups of order p_n , one subgroup of order q_n , and one subgroup of order $p_n q_n$. Then

$$\sigma_1(G_n) = \frac{1 + p_n q_n + q_n + p_n q_n}{p_n q_n} = 2 + \frac{1 + \frac{1}{q_n}}{p_n},$$

and clearly $\sigma_1(G_n) \searrow 2$ for $n \rightarrow \infty$, as desired. $\square$

References

- [1] T. De Medts and M. Tărnăuceanu, *Finite groups determined by an inequality of the orders of their subgroups*, Bull. Belg. Math. Soc. Simon Stevin **15** (2008), 699-704.
- [2] R. Schmidt, *Subgroup lattices of groups*, de Gruyter Expositions in Mathematics 14, de Gruyter, Berlin, 1994.
- [3] M. Suzuki, *Group theory*, I, II, Springer-Verlag, Berlin, 1982, 1986.
- [4] M. Tărnăuceanu, *Groups determined by posets of subgroups*, Ed. Matrix Rom, București, 2006.

Marius Tărnăuceanu
Faculty of Mathematics
‘‘Al.I. Cuza’’ University
Iași, Romania
e-mail: `tarnauc@uaic.ro`